FORM PTO-1449/A and B (Modified)	APPLICATION NO.: 09/913,756	DOCKET NO.: L0461.70121US00	
INFORMAZION DISCLOSURE	INTL. FILING DATE: 02/18/00	Confirmation No.: 5298	
STATEMENTARY APPLICANT	APPLICANT: Chiari, et al.		
Sheet 1 of 3	GROUP ART UNIT: 1644	EXAMINER: Vandervegt, F.	

U.S. PATENT DOCUMENTS

Examiner's Initials	Cite No.	U.S. Patent Docum	Kind Code	Name of Patentee or Applicant of Cited Document	Date of Publication or of issue of Cited Document MM-DD-YYYY
R/	Al	5,342,774	Ì	Boon et al.	08-30-1994

FOREIGN PATENT DOCUMENTS

Examiner's Cite Foreign Patent Document		Name of Patentee or Applicant of Cited Document	Date of Publication of	Translation			
Initials	No.	Office/ Country	Number	Kind Code	(not necessary)	Cited Document MM-DD-YYYY	(Y/N)
R	Bl	wo	93/00425	A1	The Walter and Eliza Hall Institute of Medical Research	07-01-1993	
	B2	wo	95/25740	Al	Ludwig Institute for Cancer Research et al.	09-28-1995	
	B3	wo	97/11669	A2	The Government of the United States of America	04-03-1997	<u></u>
 	B4	wo	97/31017	Al	Ludwig Institute for Cancer Research et al.	08-28-1997	
<u>'</u>	B5	wo	99/14326	Al	Ludwig Institute for Cancer Research et al.	03-25-1999	

OTHER ART — NON PATENT LITERATURE DOCUMENTS

		OTHER ART — NON PATENT LITERATURE DOCUMENTS	Translation
Examiner's	er's Cite Include name of the author (in CAPITAL LETTERS) title of the article (when appropriate), title of the item		(Y/N)
Initials	No	(book, magazine, journal, serial, symposium, catalog, etc.), date, relevant page(s), volume-issue number(s),	
		publisher, city and/or country where published.	
R/	G.	ANDERSON et al., Unified nomenclature for Eph family receptors and their ligands, the ephrins.	
VY	Cl	Enh Namonalatura Committee Cell 1997 Aug 8:90(3):403-4.	
	<u></u>	AVANZI et al. M-07e human leukemic factor-dependent cell line provides a rapid and sensitive	
	C2	L biography for the human cytokines GM-CSF and IL-3, I Cell Physiol, 1990 Dec; 145(3):458-04.	
	C3	ROYD et al. Isolation and characterization of a novel receptor-type protein tyrosine kinase (nek)	l .
i i	C3	from a human pre-B cell line. J Biol Chem. 1992 Feb 15;267(5):3262-7. (XP-000615518)	Ļ
 		BRICHARD et al. A tyrosinase nonapeptide presented by HLA-B44 is recognized on a human	.
.	C4	I melanoma by autologous cytolytic T lymphocytes. Eur J Immunol. 1996 Jan;20(1):224-30.	
		CHAUX et al., Identification of MAGE-3 epitopes presented by HLA-DR molecules to CD4(+) T	2.555
ł	C5	L	
 	CHIARLet al. Identification of a tumor-specific shared antigen derived from an Eph receptor		
presented to CD4 T cells on HLA class II molecules. Cancer Res. 2000 Sep 1;60		presented to CD4 T cells on HLA class II molecules. Cancer Res. 2000 Sep 1;60(17):4853-63.	
		CHICZ et al., Specificity and promiscuity among naturally processed peptides bound to HLA-DR	
1	C7	ollolog Evp Med 1993 Jul 1:178(1):27-47	
	 	CONNOR et al., Genomic organization and alternatively processed forms of Ceks, a receptor	ļ
į	C8	protein-tyrosine kinase of the Enh subfamily. Oncogene, 1995 Dec /:11(11):2429-38.	
		COULIE et al., Antigens recognized on human tumors by cytolytic T lymphocytes: towards	į
1	C9		
		DE PLAEN et al., Structure, chromosomal localization, and expression of 12 genes of the MAGE	
i i	C10	family Immunogenetics, 1994:40(5):360-9. (SP000614537)	<u></u>
		DOTTORI et al. Cloning and characterization of EphA3 (Hek) gene promoter: DNA methylation	l
\	CII	regulates expression in hematopoietic tumor cells. Blood. 1999 Oct 1;94(7):2477-86. (XP-	
1	J C11,	000907581)	
	-	ENGELHARD, Structure of peptides associated with class I and class II MHC molecules. Annu	
 `	C12	Box Immunol 1004:12:181.207	1
	-	Rev Immunol. 1994;12:181-207. GILBERT et al., A protein particle vaccine containing multiple malaria epitopes. Nat Biotechnol.	1
1	C13		
	ــــــــــــــــــــــــــــــــــــــ	1997 Nov;15(12):1280-4	

0 d. franklin 2/3/05

790840.1

PADEMARK

FORM PTO-1449/A and B (Modified)	APPLICATION NO.: 09/913,756	DOCKET NO.: L0461.70121US00
DEFORMATION DISCLOSURE	IN'TL. FILING DATE: 02/18/00	Confirmation No.: 5298
STATEMENT BY APPLICANT	APPLICANT: Chiari, et al.	
JAN 0 6 2005 55)	GROUP ART UNIT: 1644	EXAMINER: Vandervegt, F.
Shear 2.5 of 3		

(A)	EMARKON		
Examiner's	Cite	Include name of the author (in CAPITAL LETTERS) title of the article (when appropriate), title of the item	Translation
Initials	, No	(book, magazine, journal, serial, symposium, catalog, etc.), date, relevant page(s), volume-issue number(s), publisher, city and/or country where published.	(Y/N)
0.7	 	HEIDECKER et al., Cytolytic T lymphocytes raised against a human bladder carcinoma recognize	
001/	CI4	an antigen encoded by gene MAGE-A12. J Immunol. 2000 Jun 1;164(11):6041-5. (XP002173646)	
		HERMAN et al., A peptide encoded by the human MAGE3 gene and presented by HLA-B44	
	C15	induces cytolytic T lymphocytes that recognize tumor cells expressing MAGE3. Immunogenetics.	1
! !		1996;43(6):377-83.	
 	 	LACKMANN et al., Distinct subdomains of the EphA3 receptor mediate ligand binding and	
	C16	receptor dimerization. J Biol Chem. 1998 Aug 7;273(32):20228-37. (XP-00914515)	
} -		LALLY et al., Unmasking cryptic epitopes after loss of immunodominant tumor antigen	
l \	١,	expression. IN Immunology 2000: The American Association of Immunologists and Clinical	
]]	C17	Immunology Society Joint annual meeting. Seattle, Washington, USA. May 12-16, 2000.	
		Immunology Society Joint annual meeting. Seattle, washington, O.S.A. May 12-10, 2000.	
		Abstracts. FASEB J. 2000 Apr 20;14(6): A1005, Abstract No. 54.4. (XP002173648)	
1 .		LEHMANN et al., Differences in the antigens recognized by cytolytic T cells on two successive	
l [C18	metastases of a melanoma patient are consistent with immune selection. Eur J Immunol. 1995	
	<u> </u>	Feb;25(2):340-7.	
	C19	LI et al., IL-1 beta alters the expression of the receptor tyrosine kinase gene r-EphA3 in neonatal	
]	_	rat cardiomyocytes. Am J Physiol. 1998 Jan;274(1 Pt 2):H331-41. (XP-00913942)	
		MEAZZA et al., Interleukin (IL)-15 induces survival and proliferation of the growth factor-	
	C20	dependent acute myeloid leukemia M-07e through the IL-2 receptor beta/gamma. Int J Cancer.	
`		1998 Oct 5:78(2):189-95.	
 	—	NAKANO et al., Positive selection of T cells induced by viral delivery of neopeptides to the	
1	C21	thymus Science 1997 Ian 31:275(5300):678-83.	
	PANELLI et al. A tumor-infiltrating lymphocyte from a melanoma metastasis with decrease		
	C22	expression of melanoma differentiation antigens recognizes MAGE-12. J Immunol. 2000 Apr	
	000	15;164(8):4382-92. (XP002173647)	
 	 	PARKER et al, Scheme for ranking potential HLA-A2 binding peptides based on independent	
1	C23	binding of individual pentide side-chains, J Immunol, 1994 Jan 1:152(1):163-75.	
 		PIEPER et al., Biochemical identification of a mutated human melanoma antigen recognized by	
·	C24	CD4(+) T cells I Exp Med, 1999 Mar 1:189(5):757-66.	
		RAMMENSEE et al., MHC ligands and peptide motifs: first listing. Immunogenetics.	·
i .	C25	1995;41(4):178-228. (XP000673045)	
		SAJJADI et al., Identification of a new eph-related receptor tyrosine kinase gene from mouse and	
	C26	chicken that is developmentally regulated and encodes at least two forms of the receptor. New	
<u> </u>	020	Biol. 1991 Aug;3(8):769-78. (XP-00920929)	
	 	SANDERSON et al., Expression of endogenous peptide-major histocompatibility complex class II	
	C27	complexes derived from invariant chain-antigen fusion proteins. Proc Natl Acad Sci U S A. 1995	
	C21	Aug 1;92(16):7217-21.	
	ļ <u></u>	STEIMLE et al., Complementation cloning of an MHC class II transactivator mutated in hereditary	
l ,	Ç28	MHC class II deficiency (or bare lymphocyte syndrome), Cell. 1993 Oct 8:75(1):135-46.	
 	 `	TAM et al., Incorporation of T and B epitopes of the circumsporozoite protein in a chemically	
.	C29	defined synthetic vaccine against malaria. J Exp Med. 1990 Jan 1:171(1):299-306.	
 	 	TANG et al., A variant transcript encoding an isoform of the human protein tyrosine kinase	
	C20	EPHB2 is generated by alternative splicing and alternative use of polyadenylation signals.	{
	C30		i
 	ļ <u> </u>	Oncogene. 1998 Jul 30;17(4):521-6.	
1 1		THOMSON et al., Targeting a polyepitope protein incorporating multiple class II-restricted viral	
1	C31	epitopes to the secretory/endocytic pathway facilitates immune recognition by CD4+ cytotoxic T	
	1	lymphocytes; a novel approach to vaccine design. J Virol. 1998 Mar;72(3):2246-52.	<u> </u>

D. Rught 2/3/05

790840.1

FORM PTO-1449/A and B (Modified)	APPLICATION NO.: 09/913,756	DOCKET NO.: L0461.70121US00
INFORMATION DISCLOSURE STATEMENT BY APPLICANT	IN'TL. FILING DATE: 02/18/00	Confirmation No.: 5298
STATEMENT BY APPLICANT	APPLICANT: Chiari, et al.	
JAN 0 6 2005 &	GROUP ART UNIT: 1644	EXAMINER: Vandervegt, F.
Sheet 3 Supf 3		1

	A MARIENT	Include name of the author (in CAPITAL LETTERS) title of the article (when appropriate), title of the item	Translation
Examiner's Initials	No	publisher, city and/or country where published.	(Y/N)
R	C32	THOMSON et al., Minimal epitopes expressed in a recombinant polyepitope protein are processed and presented to CD8+ cytotoxic T cells: implications for vaccine design. Proc Natl Acad Sci U S A. 1995 Jun 20;92(13):5845-9.	
	C33	THOMSON et al., Recombinant polyepitope vaccines for the delivery of multiple CD8 cytotoxic T cell epitopes. J Immunol. 1996 Jul 15;157(2):822-6.	
	C34	TOPALIAN et al., Melanoma-specific CD4+ T cells recognize nonmutated HLA-DR-restricted tyrosinase epitopes, J Exp Med. 1996 May 1;183(5):1965-71.	
	. C35	TOPALIAN, MHC class II restricted tumor antigens and the role of CD4+ T cells in cancer immunotherapy. Curr Opin Immunol. 1994 Oct;6(5):741-5.	
	C36	TRAVERSARI et al., Transfection and expression of a gene coding for a human melanoma antigen recognized by autologous cytolytic T lymphocytes. Immunogenetics. 1992;35(3):145-52.	
	C37	VAN DEN EYNDE et al., New tumor antigens recognized by T cells. Curr Opin Immunol. 1995	
	C38	VAN DEN EYNDE et al. A new family of genes coding for an antigen recognized by autologous	
	C39	VAN DER BRUGGEN et al., A gene encoding an antigen recognized by cytolytic 1 lymphocytes on a human melanoma. Science. 1991 Dec 13;254(5038):1643-7.	
÷	Ç40	VAN DER BRUGGEN et al., A peptide encoded by human gene MAGE-3 and presented by HLA-A2 induces cytolytic T lymphocytes that recognize tumor cells expressing MAGE-3. Eur J Immunol. 1994 Dec;24(12):3038-43.	
·	C41	WANG et al., Cloning genes encoding MHC class II-restricted antigens: mutated CDC27 as a	
	C42	WICKS et al., Molecular cloning of HEK, the gene encoding a receptor tyrosine kinase expressed by human lymphoid tumor cell lines. Proc Natl Acad Sci U S A. 1992 Mar 1;89(5):1611-5. (XP-000615502)	
	C43	WU et al., Engineering an intracellular pathway for major histocompatibility complex class II presentation of antigens. Proc Natl Acad Sci U S A. 1995 Dec 5;92(25):11671-5.	
	C44	YEE et al., Isolation of tyrosinase-specific CD8+ and CD4+ T cell clones from the peripheral blood of melanoma patients following in vitro stimulation with recombinant vaccinia virus. J Immunol. 1996 Nov 1:157(9):4079-86.	
	C45	ZISCH et al., Complex formation between EphB2 and Src requires phosphorylation of tyrosine 611 in the EphB2 juxtamembrane region. Oncogene. 1998 May;16(20):2657-70. (XP-000913940)	
	C46	GENBANK Submission; NIH/NCBI; Accession number M83941; Wicks et al.; 31 December 1994 (Last Submission).	

EXAMINER:	DATE CONSIDERED:
17/1/2010/	2/3/05
Ur. The why	<u> </u>

#EXAMINER: Initial if reference considered, whether or not citation is in conformance with MPEP 609; Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to Applicant.

[NOTE - The Office hereby waives the requirement under 37 CFR 1.98 (a)(2)(i) for submitting a copy of each cited U.S. patent and each U.S. patent application publication for all U.S. national patent applications filed after June 30, 2003 and for all international applications that have entered the national stage under 35 USC 371 after June 30, 2003. See 37 CFR 1.491(b). For all patent applications filed on or before June 30, 2003, copies of cited U.S. patents and patent application publications are still required unless an eIDS is filed. Copies of all other patent(s), publication(s), or other information listed must still be provided, even if it was previously submitted to, or cited by, the U.S. Patent Office in an earlier application, unless the earlier application is identified by the IDS and is relied upon for an earlier filing date under 35 U.S.C. §120, and the copy was provided in the earlier application.]